

Dosing Instructions for hCG (Novarel, Pregnyl, Human Chorionic Gonadotropin)

<p style="text-align: center;"><u>MEDICATION DOSE: 10,000 units hCG:</u></p> <p>* IF YOU HAVE ONE 10,000IU VIAL:</p> <ol style="list-style-type: none"> 1. Draw out 1cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ENTIRE 1cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>* IF YOU HAVE TWO 5,000IU VIALS:</p> <ol style="list-style-type: none"> 1. Draw out 1cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into first powder vial. 2. Once medication has dissolved completely draw out ENTIRE 1cc/mL of mixture. 3. Inject mixture into second powder vial. Once medication has dissolved completely draw out ENTIRE 1cc/mL of mixture. 4. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. 	<p style="text-align: center;"><u>MEDICATION DOSE: 5,000 units hCG:</u></p> <p>* IF YOU HAVE ONE 10,000IU VIAL:</p> <ol style="list-style-type: none"> 1. Draw out 2cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ONLY 1cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>* IF YOU HAVE TWO 5,000IU VIALS:</p> <ol style="list-style-type: none"> 1. Draw out 1cc/mL of water and inject into powder vial. 2. Once medication has dissolved completely draw out ENTIRE 1cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>**please note- you will not use the second vial of hcg powder**</p>
<p style="text-align: center;"><u>MEDICATION DOSE: 2,500 units hCG:</u></p> <p>* IF YOU HAVE ONE 10,000IU VIAL:</p> <ol style="list-style-type: none"> 1. Draw out 2cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ONLY 0.5cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>* IF YOU HAVE TWO 5,000IU VIALS:</p> <ol style="list-style-type: none"> 1. Draw out 1cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ONLY 0.5cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>**please note- you will not use the second vial of hcg powder**</p>	<p style="text-align: center;"><u>MEDICATION DOSE: 1,500 UNITS hCG:</u></p> <p>* IF YOU HAVE ONE 10,000IU VIAL:</p> <ol style="list-style-type: none"> 1. Draw out 3cc/mL of water using the 3ml 18 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ONLY 0.5cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>* IF YOU HAVE TWO 5,000IU VIALS:</p> <ol style="list-style-type: none"> 1. Draw out 1cc/mL of water using the 3ml 18g 1 ½” needle/syringe and inject into powder vial. 2. Once medication has dissolved completely draw out ONLY 0.3cc/mL of mixture. 3. Change to a 30g ½” or 27g ½” needle and inject to the lower abdomen or outer thigh at your assigned time. <p>**please note- you will not use the second vial of hcg powder**</p>
<p style="text-align: center;"><u>LUPRON TRIGGER INSTRUCTIONS</u></p> <p>* If your Lupron has not been drawn up for you:</p> <ol style="list-style-type: none"> 1. Draw out 40 units of Lupron from vial using the provided orange capped insulin syringes 2. Inject medication anywhere on the abdomen below the belly button <p>* If your Lupron has already been drawn up for you:</p> <ol style="list-style-type: none"> 1. Inject medication anywhere on the abdomen below the belly button line using the provided orange capped insulin syringe 	<p>**Please note – you may be instructed to inject your hCG intramuscularly**</p> <p>If so, you will need to change to the 22g 1 ½’ needle, and inject your dose to the UPPER OUTER QUADRANT OF YOUR BUTTOCKS at your assigned time.</p>